

## *Nervilia cumberlegei* (Orchidaceae), a Newly Recorded Orchid from Myanmar

Myo Min Latt<sup>1,2</sup>, Byung Bae Park<sup>1</sup> and Nobuyuki Tanaka<sup>3,\*</sup>

<sup>1</sup>Department of Environment and Forest Resources, College of Agriculture and Life Science, Chungnam National University, Republic of Korea

<sup>2</sup>Department of Environmental Economic, Policy and Management, University of Forestry and Environmental Sciences, Yezin, Myanmar

<sup>3</sup>Department of Botany, National Museum of Nature and Science, 4–1–1 Amakubo, Tsukuba, Ibaraki 305–0005, Japan

\*E-mail: nobuyuki\_tanaka@kahaku.go.jp

(Received 6 November 2020; accepted 23 December 2020)

**Abstract** *Nervilia cumberlegei* (Orchidaceae) is recorded in Myanmar for the first time. A description, locality information and photographs are provided. Thus far *N. cumberlegei* is known only from Taiwan and Thailand. This discovery in Myanmar represents the western edge of the species distribution.

**Key words** : Burma, section *Linervia*, Tanintharyi, taxonomy, terrestrial orchid.

### Introduction

*Nervilia* Comm. ex Gaudich. is commonly known as ‘shield orchid’. Most species grow in forested habitats and are shade-demanders (Gale *et al.*, 2018). The genus *Nervilia* is most diverse in tropical Asia, although it is widely distributed in the Old World tropics, from Australasia and the South-west Pacific Islands to sub-Saharan Africa (Pettersson, 1991; Gale *et al.*, 2015, 2018). There are approximately 71 species in the genus (Tang *et al.*, 2018; WCSP, 2020). However, *Nervilia* remains poorly known, especially in the Asian region (Gale *et al.*, 2014). This genus is troublesome in taxonomic identification because of the development of flower and leaves in different seasons (Chen and Gale, 2009). The populations are often small, isolated and prone to demographic stochasticity (Gale *et al.*, 2010; Gale *et al.*, 2018). *Nervilia* is characterized by having a prominent fan-shaped leaf venation

(Pridgeon *et al.*, 2005; Tang *et al.*, 2018) and flowers followed by a vegetative stem bearing a single photosynthetic leaf (Niissalo *et al.*, 2020). *Nervilia* plants have antioxidant and antibacterial properties (Ruthisha *et al.*, 2018) and are used as traditional medicine. Medicinal orchids are considerably threatened by anthropogenic impacts such as habitat destruction, and illegal collection for commercial and subsistence uses (Pant *et al.*, 2002; Pant, 2013).

In Myanmar, Kress *et al.* (2003) and Aung *et al.* (2020) recorded six species of *Nervilia* in their checklists; *Nervilia concolor* (Blume) Schltr. (= *N. aragoana* Gaudich.), *N. juliana* (Roxb.) Schltr., *N. khasiana* (King & Pantl.) Schltr., *N. macroglossa* (Hook.f.) Schltr., *N. maculata* (C.S.P.Parish & Rchb.f.) Schltr. [= *N. parishiana* (King & Pantl.) Schltr.] and *N. plicata* (Andrews) Schltr. In the latest checklist by Aung *et al.* (2020), the number of species has not been changed, however, some names are updated. During a field trip to Tanintharyi Region, *Nervilia cumberlegei* Seidenf. & Smiti-




Fig. 1. *Nervilia cumberlegei* and its habitat. A. Habitat. B. Leaf. C. Tubers. D. Inflorescence.

nand was collected, which was previously known only from Taiwan and Thailand. This is the first record of *N. cumberlegei* from Myanmar.

### Materials and Methods

The field survey in Tanintharyi Nature Reserve and the adjacent area around San-daw-shin Pagoda Hill, was carried out from April to August 2009. Herbarium specimens and living material were collected and photographs were taken. The measurements were taken in the field. Voucher specimens are kept in the Education Center of Tanintharyi Nature Reserve and in the herbarium of the Forest Research Institute, Myanmar (RAF). The living specimens are cultivated at the Orchid House of Tanintharyi Nature Reserve and are used for study.

### Results and Discussion

The study area, ranging up to 10 km from the coastal line to the intermittent mountain range in Tanintharyi Nature Reserve, is covered by dry deciduous forest dominated by *Dipterocarpus obtusifolius* Teijsm. ex Miq. (Dipterocarpaceae). In April 2009, two *Nervilia* inflorescences were collected in bamboo leaf litter near Zinbar stream. From May to August 2009, several plants with leaves were discovered along Zinbar stream to Heinze stream as well as near San-daw-shin Pagoda. The *Nervilia* plants did not match any descriptions of the six species recorded in the latest checklist (Aung *et al.*, 2020). After careful examination of both herbarium specimen and living material, it was identified as *Nervilia cumberlegei* Seidenf. & Smitinand, described from Khao Yai National Park, central Thailand. It is characterized by having an 2–4-flowered inflo-

rescence and green flowers with white and strongly fimbriate labellum.

### Taxonomic Treatment

***Nervilia cumberlegei*** Seidenf. & Smitinand, Orch. Thail. (Prelim. List): 729 (1965); Seidenf. in Dansk Bot. Ark. 32(2): 155 (1978); Leou in Quart. J. Expt. Forest Natl. Taiwan Univ. 11(2): 75 (1997); Su in Editorial Committee of the Flora of Taiwan (ed.), Fl. Taiwan, ed. 2, 5: 978 (2000); Chen and Gale, in Wu *et al.*, (eds), Fl. China 25: 200 (2009); Gale and Wathana in Balslev *et al.* (eds), Fl. Thailand 12(2): 559 (2014).

[Fig.1 B–D]

Terrestrial herb growing among leaf litter, flowering when leafless, up to 8 cm tall in flower, leaf adpressed to the ground. **Tuber** white, globose, 8 × 10 mm. **Subterranean stem** from the apical node of the tuber, 2.7 cm long, 3 mm in diam. **Leaf blade** 4–5 cm in diam., dark green, reniform, fan-shaped, glabrous, with 9 veins, thick and leathery, developing after flowering. **Inflorescence** 8 cm long, 3–4-flowered; peduncle 1.5 cm long, 3 mm in diam., smooth, green, bearing linear-lanceolate, pale green, tubular sheaths, sheaths 4 × 20 mm. **Flowers** green with white fringed labellum, 3 cm across, sepals and petals free, pedicel short, 2 mm long, green; **sepals** green, narrowly elliptic, apex acute; dorsal sepal 15 × 2 mm; lateral sepals 15 × 2 mm; petal narrowly elliptic, 14.5 × 2 mm; **labellum** white, 3-lobed, oblong, 15 × 4 cm, lateral lobes round, mid-lobe fringed, pubescent; disk white with green veins. **Column** white, erect, ‘hugged’ by lateral lobes of labellum, 10 mm long, hypochile greenish; anther cap white. **Ovary** conspicuous, green, 3-angled, 4 mm long.

*Specimens examined*: —MYANMAR. Tanintharyi Region. New Mi-chaung-laung and Heinze Local Operation Unit, Tanintharyi Nature Reserve, Yephyu Township, Dawei District, ca. 252 m elev., 14°30′03.6″N, 98°13′17.9″E, 25 April 2009, *Myo Min Latt 003* (RAF).

*Habitat*: —*Nervilia cumberlegei* grows in a humus layer over clay in bamboo forest and in stony soil in dry deciduous forest (Fig. 1A).

*Phenology*: —Flowering in April to May before the rainy season.

*Distribution*: —Myanmar (new record), Taiwan and Thailand.

*Note*: —We propose the local name of this species as “Tanintharyi Ta-bin-shwe-htee”. Ta-bin-shwe-htee is the Myanmar name of *Nervilia*. *Nervilia cumberlegei* was previously known only from Taiwan and Thailand. The discovery in Myanmar represents the western limit of the species distribution (Fig. 2).

*IUCN Conservation Status*: —Many populations were found in dry deciduous forest dominated by *Dipterocarpus obtusifolius*. However, the extent of the populations remains unknown, and the habitat is very close to rubber plantations and human settlements. The species may be threatened by further expansion of the rubber plantations and other agricultural land development. Therefore *N. cumberlegei* is here assessed as Near Threatened (NT) (IUCN, 2012).




Fig. 2. The locality of *Nervilia cumberlegei* in Myanmar. A red circle indicates the newly recorded population in Myanmar, and black circles known populations in Khao Yai National Park, Thailand, and near Sun Moon Lake, central Taiwan, based on Seidenfaden and Smitinand (1965), and Su (2000).

## Acknowledgments

This study was carried out by the University of Forestry and Environmental Science (UFES), Yezin, and by the biodiversity conservation program of Tanintharyi Nature Reserve Project (TNRP) which is the first collaborative project between the Ministry of Environmental Conservation and Forestry (presently Ministry of Natural Resources and Environmental Conservation), Union of Myanmar, and PETRONAS Company together with Total E & P Myanmar. The authors are grateful to the Forest Department of Myanmar for permitting and supporting our botanical surveys in the protected areas. We sincerely thank TNRP staff Bo Bo Soe and Kyaw Thura for their assistance in the field.

## References

- Aung, Y. L., Mu, A. T., Aung, M. H., Liu, Q. and Jin, X.-H. 2020. An annotated checklist of Myanmar orchid flora. *PhytoKeys* 138: 49–112.
- Chen, X. and Gale, S. W. 2009. Orchidaceae. In: Wu, Z. Y. and Raven, P. H. (eds), *Flora of China* vol. 25, pp. 197–201. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis.
- Gale, S. W., Rueangrua, S. and Suddee, S. 2014. Studies in Asian *Nervilia* (Nervilieae, Epidendroideae, Orchidaceae) IV: *N. umphangensis*, a new species from the Thai–Myanmar border. *Phytotaxa* 166(2): 139–144.
- Gale, S. W., Li, J., Kinoshita, A. and Yukawa, T. 2015. Studies in Asian *Nervilia* (Orchidaceae) V: *N. futago*, a cryptic new species from southwest Japan confirmed by morphological, cytological and molecular analyses. *Systematic Botany* 40: 413–425.
- Gale, S. W., Maeda, A., Chen, C. I. and Yukawa, T. 2010. Interspecific relationships and hierarchical spatial genetic structuring in *Nervilia nipponica*, an endangered orchid in Japan. *Journal of Plant Research* 123: 625–637.
- Gale, S. W., Duangjai, S., Li, J., Ito, Y., Watthana, S., Termwutthipreecha, P., Cheuk, M. L. and Suddee, S. 2018. Integrative analyses of *Nervilia* (Orchidaceae) section *Linervia* reveal further undescribed cryptic diversity in Thailand. *Systematics and Biodiversity* 16(4): 377–396. DOI: 10.1080/14772000.2017.1415233
- IUCN. 2012. Red List Categories and Criteria: ver. 3.1. 2nd ed. IUCN, Gland, Switzerland and Cambridge, UK.
- Kress, W. J., DeFilipps, R. A., Farr, E. and Kyi, Y. Y. 2003. A checklist of the trees, shrubs, herbs, and climbers of Myanmar. *Contributions from the United States National Herbarium* 45: 118–124.
- Niissalo, M. A., Choo, L. M., Kurzweil, H., Yam, T. W. and Khew, G. S. 2020. A new species of *Nervilia* (Orchidaceae) from Singapore. *Gardens' Bulletin Singapore* 72(1): 1–14. doi: 10.26492/gbs72(1).2020-01.
- Pant, B., Chaudhary, R. P., Subedi, A. and Shakya, L. R. 2002. Nepalese Himalayan Orchids and the conservation priorities. In: *Proceeding of International Seminar on Mountains*, pp. 485–495. Royal Nepal Academy of Science and Technology.
- Pant, B. 2013. Review: Medicinal orchids and their uses: Tissue culture a potential alternative for conservation. *African Journal of Plant Science* 7(10): 448–467. DOI: 10.5897/AJPS2013.1031
- Pettersson, B. 1991. The genus *Nervilia* (Orchidaceae) in Africa and the Arabian Peninsula. *Orchid Monographs*, Volume 5. Leiden: Rijksherbarium/Hortus Botanicus.
- Pridgeon, A. M., Cribb, P. J., Chase, M. W. and Rasmussen, F. N. (eds). 2005. *Genera Orchidacearum*, vol. 4, Epidendroideae, pt. 1. Oxford: Oxford University Press.
- Ruthisha P. K., Krishna, S. C. and Khaleel, K. M. 2018. Phytochemical, antioxidant, antibacterial activities of *Nervilia crocifformis* (Zoll. & Moritzii) Seidenf. and *Nervilia infundibulifolia* Blatt. & McCann (Orchidaceae) collected from Kasargod District, Kerala, India. *International Journal of Pharmaceutical Science and Research* 9(12): 5177–5185.
- Seidenfaden, G. and Smitinand, T. 1965. The orchids of Thailand: a preliminary list. Part IV(2). The Siam Society, Bangkok.
- Su, H. J. 2000. Orchidaceae. In: *Editorial Committee of the Flora of Taiwan* (ed.), *Flora of Taiwan*, ed. 2, vol. 5, pp. 978–1088. National Taiwan University, Taipei.
- Tang, F.-X., Li, J.-W., Pan, B., Wu, X.-F., Luo, Y. and Liu, Q. 2018. New and Newly Record Orchids of *Nervilia* (Nervilieae, Epidendroideae, Orchidaceae) in China. *Phytotaxa* 379 (2): 162–168.
- WCSP. 2020. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. <http://wesp.science.kew.org/> [Accessed 9 December 2020].