

## Karyotypes of Four Cyprinid Fishes from Thailand

By

**Ryoichi ARAI**

Department of Zoology, National Science Museum, Tokyo

and

**Wichian MAGTOON**

Department of Biology, Faculty of Science, Srinakharinwirot University  
at Bangkok, Bangkok, Thailand

**Abstract** Chromosomes of four cyprinid fishes from Thailand were observed. Karyotypes of these four species are as follows: *Mystacoleucus marginatus*:  $2n=50$ ,  $NF=76$ ; *Acrossocheilus deauratus*:  $2n=50$ ,  $NF=72$ ; *Puntius orphoides*:  $2n=50$ ,  $NF=80$ ; *Labeo behri*:  $2n=50$ ,  $NF=70$ . Classification of *Acrossocheilus* is discussed from karyological and morphological viewpoints. The karyotype of *Mystacoleucus marginatus* is the first reported in the genus *Mystacoleucus*.

In order to clarify the phylogenetic systematics and zoogeography of cyprinid fishes in Thailand, we have observed karyotypes of cyprinid fishes from Thailand (MAGTOON & ARAI, 1989 and 1990). Recently, we had the chance to examine chromosomes of 4 cyprinid fishes in Thailand, i. e., *Mystacoleucus marginatus* (VALENCIENNES in CUVIER & VALENCIENNES, 1842), *Acrossocheilus deauratus* (VALENCIENNES in CUVIER & VALENCIENNES, 1842), *Puntius orphoides* (VALENCIENNES in CUVIER & VALENCIENNES, 1842), and *Labeo behri* FOWLER, 1937.

As karyotypes of these four species have not been reported, the results will be described shortly in the following lines.

### Materials and Method

Three specimens of *Labeo behri*, 86.5 to 93.6 mm in total length (TL), and a specimen of *Acrossocheilus deauratus*, 133.8 mm in TL, were collected from Ubon Ratchathani, Northeast Thailand. A specimen of *Puntius orphoides*, 95.9 mm in TL, was caught at Mae Hong Song, North Thailand, and two specimens of *Mystacoleucus marginatus*, 87.2 and 102.7 mm in TL, at Uthai Thani, Central Thailand. Some morphological characters of material fishes are shown in Table 1. They are deposited in the fish collection of the Department of Zoology, National Science Museum, Tokyo. The classification of the family Cyprinidae in Thailand follows that by SMITH (1945).

Table 1. Morphological characters of material fishes.

Species	No. of fish	Standard length (mm)	Barbels	Lateral line scales*	Branched dorsal rays	Branched anal rays	Vertebrae
<i>Mystacoleucus marginatus</i>	2	65.5–76.5	4	27–30(2)	8	9	34
<i>Acrossocheilus deauratus</i>	1	104.6	4	31(3)	8	5	38
<i>Puntius orphoides</i>	1	73.6	4	28(2)	8	5	33
<i>Labeo behri</i>	3	62.2–73.8	2	39(3)	12	5	35

\* Scales on the body side and the caudal fin were included. Figures in parentheses show the number of scales on the caudal fin.

Chromosome preparation was made from the head kidney following the method of OJIMA and KURISHITA (1980). The classification of chromosomes is adopted from LEVAN *et al.* (1964). Metacentrics and submetacentrics are described as two-arm chromosomes, and subtelocentrics and acrocentrics as one-arm chromosomes. The definition of the new arm number is referred to ARAI and NAGAIWA (1976).

### Results

***Mystacoleucus marginatus*** (Fig. 1 A, C). As shown in Table 2, the diploid chromosome number is 50. The karyotype comprises 16 metacentric, 10 submetacentric, and 24 acrocentric chromosomes. The arm number is 76. The new arm number is 50.

***Acrossocheilus deauratus*** (Fig. 1 B, D). The diploid chromosome number is 50. The karyotype comprises 10 metacentric, 12 submetacentric, and 28 acrocentric chromosomes. The largest chromosome is submetacentric. The arm number is 72. The new arm number is 50.

***Puntius orphoides*** (Fig. 2 A, C). The diploid chromosome number is 50. The karyotype of this species comprises 14 metacentric, 16 submetacentric, 4 subtelocentric, and 16 acrocentric chromosomes. The largest chromosome is submetacentric. The arm number is 80. The new arm number is 50.

***Labeo behri*** (Fig. 2 B, D). The diploid chromosome number is 50. The karyotype comprises 12 metacentric, 8 submetacentric, 2 subtelocentric, and 28 acrocentric chromosomes. The largest chromosome is submetacentric. The arm number is 70. The new arm number is 50.

Table 2. Frequency distributions of diploid chromosome counts in four species of material fishes.

Species	2n									Total
	44	45	46	47	48	49	50	51	52	
<i>Mystacoleucus marginatus</i>			1		1		8			10
<i>Acrossocheilus deauratus</i>		1			1		6			8
<i>Puntius orphoides</i>					1		13			14
<i>Labeo behri</i>			1		1		9			11




Fig. 1. Photomicrographs of mitotic metaphase chromosomes and karyotypes of two species in the genera *Mystacoleucus* and *Acrossocheilus*. A and C, *Mystacoleucus marginatus*; B and D, *Acrossocheilus deauratus*. Each scale indicates 5  $\mu$ m.

### Discussion

Morphologically, the genus *Mystacoleucus* has the peculiar character, i.e., presence of a procumbent predorsal spine. HORA (1937) considered this character to be an autapomorphy of *Mystacoleucus*, but this character was also found in *Spinibarbus* species and *Rohtee ogilbii* (OSHIMA, 1919; SMITH, 1945). Although the karyotype of *Rohtee ogilbii* has not been reported, the diploid chromosome number of *Mystacoleucus* is 50, while that of *Spinibarbus* is 100. This fact suggests that the




Fig. 2. Photomicrographs of mitotic metaphase chromosomes and karyotypes of two species in the genera *Puntius* and *Labeo*. A and C, *Puntius orphoides*; B and D, *Labeo behri*. Each scale indicates 5  $\mu$ m.

procumbent predorsal spine is not an autapomorphy of *Mystacoleucus* and occurred independently in *Mystacoleucus*, *Spinibarbus*, and *Rohtee ogilbii*.

The karyotype of *Acrossocheilus deauratus* is the first reported in the species of this genus from Thailand. KOTTELAT (1989, tab. 3) divided *Acrossocheilus* sensu SMITH (1945) into 3 genera, i. e., *Acrossocheilus* OSHIMA, 1919, *Poropuntius* SMITH, 1931, and *Neolissochilus* RAINBOTH, 1985. When we follow the classification by

Table 3. Karyotypes of the genus *Acrossocheilus* sensu SMITH (1945).

Excepting data of present study, data sources of karyotypes are different from those of morphology. As regards arm numbers (NF), metacentrics and submetacentrics are counted as two in NF<sub>1</sub>, while metacentrics, submetacentrics, and subtelocentrics counted as two in NF<sub>2</sub>. Morphological data are adopted from DAY (1875–1878), SMITH (1945), and WU *et al.* (1977).

Species	2n	NF <sub>1</sub>	NF <sub>2</sub>	No. of scales		Vertebrae	Literature on karyotypes
				I*	II*		
<i>Acrossocheilus fasciatus</i>	50	80	90	16	37–38	37–40	GUI <i>et al.</i> , 1986
<i>A. hemispinis hemispinis</i>	50	76	84	16	36–37	37–39	YU <i>et al.</i> , 1989
<i>A. iridescens zhuijiangensis</i>	50	80	90	16	39–41		GUI <i>et al.</i> , 1986
<i>A. labiatus</i>	50	78	82	16	36–38	38–39**	ARAI, 1982
<i>A. parallens</i>	50	80	94	16	36–37		GUI <i>et al.</i> , 1986
<i>A. wenchowensis beijiangensis</i>	50	80	94	16	36–37		GUI <i>et al.</i> , 1986
<i>A. yunnanensis</i>	50	84		16–18	42–44	44	ZAN <i>et al.</i> , 1984
"	50	78	90				LI <i>et al.</i> , 1986
<i>Poropuntius deauratus</i>	50	72	72	14	29–33	38	Present study
<i>Neolissochilus hexagonolepis</i>	100	140		12	28–31	42–43**	MATSUDA, 1985
<i>N. sumatranus</i>	98	142	158	12	23–25		SUZUKI & TAKI, 1981

\* I, Number of circumpeduncular scales. II, Number of scales in lateral series.

\*\* Present study.

KOTTELAT (1989, p. 10), *Acrossocheilus deauratus* is placed in the genus *Poropuntius*. As shown in Table 3, the diploid chromosome number of *Acrossocheilus*, *Poropuntius*, and *Neolissochilus* is 50, 50, and 98 to 100, respectively. Karyologically it seems difficult to separate *Acrossocheilus* from *Poropuntius*. In case that the genus *Poropuntius* is considered to be valid, the karyotype of *P. deauratus* (= *Acrossocheilus deauratus*) is the first reported in the genus *Poropuntius*. As suggested by ARAI (1982, p. 145), *Neolissochilus* is more closely related to *Tor* than to *Acrossocheilus* sensu stricto and *Poropuntius* in the karyotype as well as morphological characters.

KOTTELAT (1989) synonymized *Labeo behri* with *Bangana pierrei*. When we follow KOTTELAT's classification, the karyotype of *B. pierrei* (= *Labeo behri*) is the first reported in the genus *Bangana*.

### Acknowledgment

We wish to express our gratitudes to Mr. Thawat DONSAKUL, Srinakharinwirot University at Bangkok, for collecting specimens of *Mystacoleucus marginatus*.

### References

- ARAI, R., 1982. A chromosome study on two cyprinid fishes, *Acrossocheilus labiatus* and *Pseudorasbora pumila pumila*, with notes on Eurasian cyprinids and their karyotypes. *Bull. natn. Sci. Mus., Tokyo*, (A), **8**: 131–152.

- ARAI, R., & K. NAGAIWA, 1976. Chromosomes of tetraodontiform fishes from Japan. *Bull. natn. Sci. Mus., Tokyo*, (A), 2: 59–72, pls. 1–6.
- DAY, F., 1875–1878. The fishes of India; being a natural history of the fishes known to inhabit the seas and fresh waters of India, Burma, and Ceylon. Vol. 1. Reprints of Today's & Tomorrow's Book Agency, xx+778 pp. New Dehli.
- GUI, J., Y. LI, K. LI & T. ZHOU, 1986. Studies on the karyotypes of Chinese cyprinid fishes. (VIII) Karyotypic analyses of fifteen species of Barbinae with a consideration for their phyletic evolution. *Trans. Chinese Ichthyol. Soc.*, 5: 119–127, pls. 1–2. (In Chinese with English abstract.)
- HORA, S. L., 1937. Systematic position, geographical distribution and evolution of the cyprinoid genera with a procumbent predorsal spine. *Rec. Ind. Mus.*, 39: 311–319.
- KOTTELAT, M., 1989. Zoogeography of the fishes from Indochinese Inland Waters with an annotated check-list. *Bull. zool. Mus., Univ. Amsterdam*, 12(1): 1–54.
- LEVAN, A., K. FREDGA & A. A. SANDBERG, 1964. Nomenclature of centromeric position on chromosomes. *Hereditas*, 52: 201–220.
- LI, Y., K. LI, J. JIANG, Q. SUN & T. ZHOU, 1986. Studies on the karyotypes of Chinese cyprinid fishes. X. Karyotypes of five species of Barbinae and four species of Gobioninae. *Zool. Res. (Kunming)*, 7: 183–189, pls. 1–3. (In Chinese with English abstract.)
- MAGTOON, W., & R. ARAI, 1989. Karyotypes of five *Puntius* species and one *Cyclocheilichthys* species (Pisces, Cyprinidae) from Thailand. *Bull. natn. Sci. Mus., Tokyo*, (A), 15: 167–175.
- & ——— 1990. Karyotypes of three cyprinid fishes, *Osteochilus hasselti*, *O. vittatus*, and *Labiobarbus lineatus*, from Thailand. *Japan. J. Ichthyol.*, 36: 483–487.
- MATSUDA, M., 1985. Chromosomes of *Acrossocheilus hexagonolepis*. *Ann. Rep. Biwakobunkakan*, 3: 1–3. (In Japanese.)
- OJIMA, Y., & A. KURISHITA, 1980. A new method to increase the number of mitotic cells in the kidney tissue for fish chromosome studies. *Proc. Japan Acad.*, 56B: 610–615.
- OSHIMA, M., 1919. Contributions to the study of the fresh water fishes of the Island of Formosa. *Ann. Carnegie Mus.*, 12: 169–328, pls. 48–53.
- RAINBOTH, W. J., 1985. *Neolissochilus*, a new genus of South Asian cyprinid fishes. *Beaufortia*, 35: 25–35.
- SMITH, H. M., 1931. Descriptions of new genera and species of Siamese fishes. *Proc. U. S. natn. Mus.*, 79 (7): 1–48, pl. 1.
- 1945. The fresh-water fishes of Siam, or Thailand. *Bull. U. S. natn. Mus.*, 188: 1–622.
- SUZUKI, A., & Y. TAKI, 1981. Karyotype of tetraploid origin in a tropical Asian cyprinid, *Acrossocheilus sumatranus*. *Japan. J. Ichthyol.*, 28: 173–176.
- WU, H. W., & 9 others, 1977. Cyprinid fishes of China, 2. Pp. 1–2+229–598, pls. 1–109. Shanghai People's Press, Shanghai. (In Chinese.)
- YU, X., T. ZHOU, Y. LI, K. LI & M. ZHOU, 1989. Chromosomes of Chinese fresh-water fishes. vi+179 pp., Science Press, Beijing. (In Chinese.)
- ZAN, R., Z. SONG & W. LIU, 1984. Studies of karyotypes of seven species of fishes in Barbinae, with a discussion on identification of fish polyploids. *Zool. Res. (Kunming)*, 5: 82–90, 2 pls. (In Chinese with English abstract.)