

A New Species of the Sawfly Genus *Masaakia* (Hymenoptera, Tenthredinidae) from Hokkaido, Japan

Ichiji Togashi

1-chome, Honmachi, Tsurugi-machi, Ishikawa Prefecture, 920–2121 Japan

Abstract *Masaakia shinoharai* sp. nov. from Hokkaido, Japan, is described and illustrated, and a key is provided for two Japanese species of *Masaakia*.

Key words: Hymenoptera, Tenthredinidae, *Masaakia shinoharai*, new species, Japan.

The genus *Masaakia* Takeuchi includes only one described species, *Masaakia longivaginata* Takeuchi, from Japan. Recently, I examined a female of this genus collected in Hokkaido, Japan. As a result of comparative study, I have concluded that this specimen represents a new species. In this paper, I describe and illustrate it.

Genus *Masaakia* Takeuchi

Masaakia Takeuchi, 1950, p. 18. Type species: *Masaakia longivaginata* Takeuchi. Monotypic.

Diagnosis. Head: occipital carina absent; clypeus with anterior margin emarginate; labrum small; malar space distinct. Antenna: 3rd segment longer than 4th (Figs. 2, 7). Thorax: prepectus absent. Hindwing without a middle cell (Figs. 3, 8). Legs: claw with a large inner tooth and basal lobe (Figs. 5, 10). Abdomen: sawsheath rather long (Figs. 11, 14).

Key to the Japanese species

1. Antenna entirely black; sawsheath as in Fig. 14; lancet as in Figs. 15–16
..... *longivaginata* Takeuchi
- Antenna with undersides of 4th to 9th segments reddish brown and apical 1/2 of underside of 3rd segment reddish brown; sawsheath as in Fig. 11; lancet as in Figs. 12–13 *shinoharai* sp. nov.


Masaakia shinoharai sp. nov.

(Figs. 1–5, 11–13)

Female. Length 6 mm. Body black, but cenchrus milky white, and cercus reddish brown. Antenna black but undersides of 4th to 9th segments and apical 1/2 of

underside of 3rd segment reddish brown (Fig. 2). Wings hyaline, stigma and veins dark brown to black. Legs black with following parts pale yellowish white: all knees, all tibiae except for dark brown apical portion, 1st segment of all tarsi except for apical portion; apical portion of 1st segment and 2nd to 5th tarsal segments dark brown to black.

Head (Fig. 1): transverse; circumocellar furrow distinct but anterior half absent; interocellar furrow short but distinct; postocellar furrow distinct (Fig. 1); lateral furrows distinct and deep; postocellar area convex, with a median longitudinal depression; OOL:POL:OCL=1.1:1.0:1.1; supra-antennal tubercles distinct; median fovea rectangular and rather shallow; lateral fovea large, circular in outline, with a small conical projection in middle; supraclypeal area nearly flattened; antenno-ocular distance slightly longer than distance between antennal sockets (ratio about 1.1:1.0); malar space long, 1/2 as long as diameter of front ocellus; postorbital groove distinct; postgenal carina distinct but rather short.



Figs. 1-5. *Masaakia shinoharai* sp. nov., holotype, ♀. — 1, Head, dorsal view; 2, antenna, lateral view; 3, wing venation; 4, inner tibial spur of fore leg, lateral view; 5, tarsal claw, lateral view. Figs. 6-10. *Masaakia longivaginata* Takeuchi, ♀, Mt. Shiritaka, Ishikawa Pref., Honshu. — 6, Head, dorsal view; 7, antenna, lateral view; 8, wing venation; 9, inner tibial spur of fore leg, lateral view; 10, tarsal claw, lateral view. Scale for 1-2 and 6-7: 1 mm; scale for 3 and 8: 5 mm; scale for 4-5 and 9-10: 0.1 mm.

Antenna slightly shorter than costa of forewing (ratio about 1.0:1.1); relative lengths of segments about 1.6:1.0:3.4:2.0:2.0:1.6:1.3:1.1:1.3; pedicel slightly longer than wide (ratio about 1.0:0.8).

Thorax: normal; mesoscutellum slightly convex. Wing venation as in Fig. 3. Legs: inner tibial spur of fore leg as in Fig. 4; hind basitarsus slightly shorter than following 4 segments combined (ratio about 1.0:1.1); tarsal claw as in Fig. 5.

Abdomen: normal; sawsheath rather long (Fig. 11); serrulae of lancet as in Figs. 12–13.


Punctuation. Vertex practically impunctate, shining; frons, supraclypeal area, clypeus, and labrum finely and sparsely punctured; temples and malar space moderately, closely, and irregularly punctured; thorax covered with very fine and sparse punctures; posterior 1/4 of mesoscutellum moderately and closely punctured; abdominal tergites practically impunctate, shining.

Male. Unknown.

Distribution. Japan (Hokkaido).

Holotype: ♀, Yamada-onsen, Tokachi, Hokkaido, 18. VI. 1991, A. Shinohara leg. Deposited in the collection of the National Science Museum (Nat. Hist.), Tokyo.

Food plant. Unknown.



Figs. 11–13. *Masaakia shinoharai* sp. nov., holotype, ♀ — 11, Sawsheath, lateral view; 12, apical portion of lancet; 13, 10th–12th serrulae of lancet. Figs. 14–16. *Masaakia longivaginata* Takeuchi, ♀, Mt. Shiritaka, Ishikawa Pref., Honshu. — 14, Sawsheath, lateral view; 15, apical portion of lancet, 16, 10th–12th serrulae of lancet. Scale for 11 and 14: 0.5 mm.

Remarks. This new species is very closely allied to *Masaakia longivaginata* Takeuchi, but it is distinguished from the latter by the coloration of the antenna (in *longivaginata*, the antenna is entirely black), by the shape of the inner tibial spur of the front leg (in *longivaginata*, the apex of the inner tibial spur is bifurcate, see Figs. 4 and 9), and by the characters of the lancet (see Figs. 12–13 and 15–16).

Acknowledgments

I express my thanks to Dr. David R. Smith, U. S. Department of Agriculture, Washington, D.C., for reading through this manuscript. I am indebted to Dr. A. Shinohara, National Science Museum (Nat. Hist.), Tokyo, for his kindness in giving me the opportunity to examine the specimen.

Reference

Takeuchi, K. 1950. An interesting new genus and species of the subfamily Blennocampinae from Japan (Tenthredinidae, Hymenoptera). *Trans. Kansai ent. Soc.*, **15** (2): 17–19.